

## Generation of a QCD-induced Chern-Simons like term in QED

D. Ebert\*, V. Ch. Zhukovsky\*<sup>†</sup>  
*Institut für Physik, Humboldt-Universität zu Berlin,  
Invalidenstraße 110, D-10115 Berlin, Germany*

### Abstract

After briefly reviewing the dynamical generation of a topological Chern-Simons (CS) term in  $QED_3$  the possibility of a Chern-Simons like term generation in the  $QED_4$  Lagrangian by quark loops interacting with a non-abelian background field of an  $SU(2)$  model of  $QCD$  is demonstrated.

---

\*Supported in part by *Graduiertenkolleg "Elementarteilchenphysik"* and *Scientific Exchange Program between Moscow State University (Moscow) and Humboldt University (Berlin)*.

<sup>†</sup>On leave of absence from the Faculty of Physics, Department of Theoretical Physics, Moscow State University, 119899, Moscow, Russia.

# 1 Introduction

In recent years much attention has been paid to the discussion of various nonperturbative effects in the theory of gauge fields, which are related to the role of topological terms and the vacuum condensate, as well as to the infrared behavior of the perturbation theory expansion at high temperature and in spaces of lower dimensions.

Some of the recently proposed QCD vacuum models are based upon investigations of the quark and gluon field interactions in the background of a certain model gluon condensate, i.e., a classical gauge field. Along with instanton models of the QCD vacuum [1], efforts are undertaken to construct models, that are based upon various regular configurations of external non-abelian fields, the simplest of which is a constant chromomagnetic field of the abelian type, proposed by Savvidy [2]. As it was shown in [3], vector-potentials, by which homogeneous and constant fields are described, can only be of two types. The first one is the so called covariantly-constant field. In this case gluon fluctuations lead to a lowering of the vacuum energy [2], while the vacuum itself turns out to be unstable [4]. The vector-potentials  $G_\mu$  of the second type are gauge equivalent to constant non-commuting potentials, so that the field tensor  $G_{\mu\nu}$  is defined through their commutator  $iG_{\mu\nu} = [G_\mu, G_\nu]$ . The chromomagnetic field configuration formed by the non-abelian potentials turned out to be stable with respect to the decay into pairs of real particles [3], [5]. In a number of publications [6], [7], modeling the vacuum by the fields of the second type, a substantially nonperturbative dependence of such objects, as the photon polarization operator, on the vacuum field was obtained. Active investigation of various radiative effects in non-abelian external fields has been initiated in this connection (see, e.g.[8]).

Since publication of the pioneer works [9], where a topological mass term for gauge fields in (2+1)-dimensional spacetime was initially investigated, effects of topology produced by Chern-Simons modification of the gauge field Lagrangian attracted much attention. The study of radiative effects in 3-dimensional theories, carried out recently in Ref. [10], [11] demonstrated the importance of the Chern-Simons (CS) topological term for the regularization of infrared divergences in calculating higher loop diagrams even in the case when background fields are present.

In particular, the one-loop electron mass operator and the photon polarization operator in (2+1)-QED in an external magnetic field at finite tem-

perature and matter density [10], as well as the mass operator of a quark in the (2+1)-dimensional QCD in an external chromomagnetic field [11] were calculated, and a nonanalytic dependence of the quantities studied on the external fields was demonstrated.

A new discussion of the role of finite matter density in the effect of the dynamical Chern-Simons term generation in (2+1)-dimensional gauge theories has been initiated recently (see e.g. [12, 13]), showing that the problem of topological effects in gauge theories is far from being solved.

A calculation of the photon polarization operator with exact consideration for the contribution of the non-abelian QCD vacuum condensate field was carried out in the case of scalar quarks in [7, 14, 15].

Of special interest are possible Lorentz- and parity-violating modifications of (3+1)-dimensional electrodynamics, investigated in [16]. These modifications are caused by adding to the *QED* Lagrangian a Chern-Simons term that couples the dual electromagnetic tensor to a four-vector. In this paper we shall address ourselves to the possibility [17] that new nontrivial topological terms of the Chern-Simons type may be generated in (3+1)-dimensional *QED*, when induced by the fermion loops in the non-abelian vacuum condensate of *QCD*.

## 2 Chern-Simons term generation in an abelian background field in (2+1) dimensions

As it is well known [9], spinor electrodynamics in (2+1)-dimensional space is described by the Lagrangian

$$\mathcal{L} = \mathcal{L}_g + \mathcal{L}_f + \mathcal{L}_{int} \quad (1)$$

with

$$\mathcal{L}_g = -\frac{1}{4}F_{\mu\nu}F^{\mu\nu} + \frac{1}{4}\theta\varepsilon^{\mu\nu\alpha}F_{\mu\nu}A_\alpha \quad (2)$$

for the gauge (electromagnetic) field, with  $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$ ,

$$\mathcal{L}_f = \bar{\psi}(i\gamma\partial - m)\psi \quad (3)$$

for the fermion field, and

$$\mathcal{L}_{int} = -j^\mu A_\mu \quad (4)$$

for the interaction term, where

$$j^\mu = -e\bar{\psi}\gamma^\mu\psi \quad (5)$$

is the fermion current. The last term in  $\mathcal{L}_g$  is the Chern-Simons term,

$$\mathcal{L}_{CS} = \frac{1}{4}\theta\varepsilon^{\mu\nu\alpha}F_{\mu\nu}A_\alpha, \quad (6)$$

related to the CS secondary characteristic class. Here  $\theta$  is the CS parameter which has a dimension of mass. It can be generated in the effective Lagrangian by quantum corrections [9, 19], which are affected by external fields, finite temperature and the density of medium (see e.g. [18, 10, 13, 12], and references therein). In the 1-loop approximation this can be seen by calculating the photon polarization operator (PO)

$$\Pi^{\mu\nu}(q) = ie^2 \int \frac{d^3p}{(2\pi)^3} \text{tr}[\gamma^\mu S(p+q/2)\gamma^\nu S(p-q/2)], \quad (7)$$

where

$$S(p) = \frac{\gamma p + m}{p^2 - m^2} \quad (8)$$

is the fermion Green's function operator. The polarization operator in  $QED_3$  has two parts

$$\Pi_{\mu\nu} = \Pi_{\mu\nu}^S + \Pi_{\mu\nu}^A, \quad (9)$$

where  $\Pi_{\mu\nu}^S$  is symmetric and  $\Pi_{\mu\nu}^A$  is antisymmetric. In the (3+1)-dimensional case  $\Pi_{\mu\nu}^A$  is absent, if there is no anisotropy in the space. It is evident that the presence of  $\Pi_{\mu\nu}^A$  means the appearance of the vacuum anisotropy, induced by vacuum corrections. The so called dynamically induced CS term in PO has the following structure:

$$\Pi_{\mu\nu}^A(q) = i\varepsilon_{\mu\nu\alpha}q^\alpha\Pi^A(q^2). \quad (10)$$

Here the value of  $\Pi^A(q^2)$  at  $q^2 = 0$  determines the induced CS topological photon mass  $\theta_{ind} = \Pi^A(0)$  [9, 19].

Note that in the presence of an external magnetic field  $H$  the function  $\Pi^A(k^2, H)$  in (10) becomes also a function of  $H$  [18, 12, 13, 10].

### 3 Chern-Simons like term generation in a non-abelian background in (3+1) dimensions

In this section we shall demonstrate the appearance of an antisymmetric structure of the type (10) in (3+1)-*QED* induced by the quark loop in a certain non-abelian background field of *QCD*. Let us consider a model with a massive quark field  $\psi$ , coupled both to an electromagnetic field  $A_\mu$  and to an  $SU(2)_C$  gluon field  $G_\mu$  in the (3+1)-dimensional space-time, which is described by the following Lagrangian

$$\mathcal{L} = \mathcal{L}_g + \mathcal{L}_f + \mathcal{L}_{int} \quad (11)$$

with

$$\mathcal{L}_g = -\frac{1}{4}F_{\mu\nu}F^{\mu\nu} - \frac{1}{4g^2}G_{\mu\nu}^a G_a^{\mu\nu}, \quad (12)$$

$$\mathcal{L}_f = \bar{\psi}(i\gamma D - m)\psi, \quad (13)$$

and  $\mathcal{L}_{int}$  is defined as in (4), (5). Here  $G_{\mu\nu}^a = \partial_\mu G_\nu^a - \partial_\nu G_\mu^a + f_{abc}G_\mu^b G_\nu^c$ , and  $D_\mu = \partial_\mu - iG_\mu$  ( $G_\mu = G_\mu^a T^a$ ,  $T^a$  are the generators of the gauge group  $SU(2)_C$ ).

If a constant non-abelian rotationally-invariant background field  $G_\mu = \text{const}$  of the second type (see Introduction) is present, then, neglecting contributions of the gluon fluctuations around this field, we may calculate the 1-loop PO of a photon ( $P_\mu = p_\mu + G_\mu$ )

$$\Pi^{\mu\nu}(q) = ie^2 \int \frac{d^4 p}{(2\pi)^4} \text{Tr} [\gamma^\mu S(P + q/2) \gamma^\nu S(P - q/2)], \quad (14)$$

moving through the QCD vacuum, and interacting with spin 1/2-quarks in the loop. Here the quark Green's function is given by the following expression with exact consideration for the vacuum field  $G_\mu$

$$S(p) = \frac{1}{\gamma P - m} = \frac{1}{P^2 - m^2 - \frac{1}{2}\sigma G}(\gamma P + m), \quad (15)$$

where  $\sigma G = \sigma_{\mu\nu} G^{\mu\nu}$ . Taking into account, that the antisymmetric term in PO appears as a structure which is proportional to an antisymmetric tensor,

we expand (15) up to a linear in  $\frac{1}{2}\sigma G$  term. For the antisymmetric part of PO

$$\Pi_{\mu\nu}^A = \frac{1}{2}(\Pi_{\mu\nu} - \Pi_{\nu\mu}), \quad (16)$$

we then obtain

$$\Pi_{\mu\nu}^A = ie^2 \int \frac{d^4 p}{(2\pi)^4} \text{Tr} \left[ \left( \frac{1}{2} \sigma G \right) \pi_{\mu\nu} \frac{1}{(P + q/2)^2 - m^2} - (\mu \leftrightarrow \nu) \right]. \quad (17)$$

Here

$$\pi^{\mu\nu}(P, q) = [\gamma^\mu S(P + q/2) \gamma^\nu S(P - q/2)]_{\sigma G=0}. \quad (18)$$

The integral in (17) is UV finite, and after further expansion in (17), (18) up to terms linear in  $G_\mu$  we obtain  $\Pi_{\mu\nu}^A$  which is nonvanishing at  $q^2 \rightarrow 0$

$$\Pi_{\mu\nu}^A(q, G) = \frac{5}{6\pi^2} \frac{e^2}{m^2} \text{tr}(G_\mu G_\nu (q_\alpha G^\alpha)). \quad (19)$$

For the rotationally-symmetric background field

$$G_1^1 = G_2^2 = G_3^3 = \sqrt{\lambda}, G_0^a = 0, \quad H_i^a = \delta_i^a \lambda (i = 1, 2, 3), \quad (20)$$

we have:

$$\Pi_{\mu\nu}^A(q, \lambda) = \frac{5i}{24\pi^2} \frac{e^2}{m^2} \lambda^{\frac{3}{2}} \varepsilon_{\mu\nu\alpha} q^\alpha, \quad (21)$$

(where now  $\mu, \nu, \alpha = 1, 2, 3$ ), which is very similar in structure to the 3d-result (10). The photon topological mass is equal to

$$\theta_{ind}(\lambda) = \frac{5}{24\pi^2} \frac{e^2}{m^2} \lambda^{\frac{3}{2}}. \quad (22)$$

We see that, like in a (2+1)-dimensional case, *QCD* vacuum corrections may induce an anisotropy in (3+1)-*QED*, if a non-abelian constant background field is present.

In the more general case of strong background fields of large intensity  $\lambda$  the induced antisymmetric structure of *QED* should have the form  $(\mu, \nu, \alpha = 1, 2, 3)$ :

$$\Pi_{\mu\nu}^A = i\varepsilon_{\mu\nu\alpha} q^\alpha \Pi^A(q^2, \lambda), \quad (23)$$

with

$$\Pi^A(0, \lambda) = \theta_{ind}(\lambda). \quad (24)$$

Using one of the eigenvectors of PO in the external field  $G_\mu$ :

$$s_\beta = -\frac{2i}{3}\lambda^{-\frac{3}{2}}\varepsilon_{\beta\sigma\lambda\rho} \operatorname{tr}(G^\sigma G^\lambda G^\rho) \quad (25)$$

we can write the corresponding CS structure in the *QED* action in a 4-dimensional covariant way

$$S_{CS} = \frac{1}{4} \int d^4x \theta_{ind}(\lambda) \varepsilon^{\mu\nu\alpha\beta} F_{\mu\nu} A_\alpha s_\beta. \quad (26)$$

One can see that the topological current vector [9], introduced for an arbitrary varying field  $G_\mu(x)$ ,

$$X^\beta = 4\varepsilon^{\beta\sigma\lambda\rho} \operatorname{tr}(G_\sigma \partial_\lambda G_\rho - \frac{2i}{3} G_\sigma G_\lambda G_\rho) \quad (27)$$

is proportional to  $s_\beta$  (25) in the case of a constant field. Thus one might expect that (26) after replacement  $s^\beta \rightarrow X^\beta$  holds also for an arbitrary varying background field with non-zero topological charge.

## 4 Discussion

The induced CS structure discussed above may provide one of the physical mechanisms for possible unusual phenomena in the propagation of light through the universe. For instance, evidence for an anisotropy in electromagnetic field propagation over cosmological distances was reported recently in [20]. They claimed to observe a new effect (extracted from Faraday rotation) of rotation of the plane of polarization of electromagnetic radiation propagating over cosmological distances, correlated with the angular position and distance of the source. It should be pointed out that this observation of anisotropy initiated many controversial comments concerning possible interpretation of the results of [20] (see e.g. [21], and a more recent publication [22] and references therein). Such an evidence, in case it is confirmed, might signal for a non-trivial topological structure of the space over large scales, explained by microscopic reasons. The authors of [20] discussed an effective

action with a gauge invariant coupling of  $A_\mu$  and  $F_{\mu\nu}$  to a certain background vector  $s_\mu$  of the form (26). The possibility that such a term in (3+1)-  $QED$  Lagrangian could produce an anisotropy in electromagnetic field propagation was pointed out first by Carroll, Field and Jackiw [16]. One might relate  $s_\mu$  to some intrinsic “spin axis” of an anisotropic universe, associated with axion type domain walls or some other specific condensate structures of the type considered above.

As it is well known, there exists a number of nontrivial topological solutions of field equations, such as, e.g., instantons. The non-abelian constant field case, discussed above, provides just a simple idea of a possible birefringence effect in the propagation of electromagnetic or other gauge fields that may arise under the influence of a nontrivial topological background. More realistic examples are to be discussed in subsequent publications.

## Acknowledgments

The authors are grateful to R.Jackiw for useful comments. One of the authors (V.Ch.Zh.) gratefully acknowledges the hospitality of Prof. Mueller-Preussker and his colleagues at the particle theory group of the Humboldt University extended to him during his stay there.

## References

- [1] A. Belavin et al., *Phys. Lett.* **B59** (1975) 85;  
E.V.Shuryak, *Nucl. Phys.* **B203** (1982) 93;  
D. I. Diakonov, M. V. Polyakov and C. Weiss, *Nucl. Phys.* **B461** (1996) 539;  
(for recent publications see A. Ringwald and F. Schrempp, “*Instanton Phenomenology at HERA*”, *Preprint DESY 97 –115*, (1997) and references therein)
- [2] G.K.Savvidy, *Phys. Lett.* **B71** (1977) 133
- [3] L.S.Brown and W.I.Weisberger, *Nucl. Phys.* **B157** (1979) 2857;
- [4] N.K.Nielsen and P.Olesen, *Nucl. Phys.* **B144** (1978) 376;



- [5] Sh.S.Agaev, A.S.Vshivtsev, V.Ch.Zhukovsky and O.F.Semyenov, *Izv.Vissht.Uch.Zav., Fizika*, **1** (1985) 78
- [6] A.I.Milshtein and Yu.F.Pinelis, *Z.Phys.* **C27** (1985) 461
- [7] A.V.Averin, A.V.Borisov and V.Ch.Zhukovsky, *Z.Phys.* **C48** (1990) 457;
- [8] I.M.Ternov, V.Ch.Zhukovsky, and A.V.Borisov, *Quantum processes in a strong external field (in Russian)*, Moscow, 1989;  
A. E. Grigoruk and V. Ch. Zhukovsky , *Vestn. Mosk. Univ., Fizika, Astron.*, **1** (1997) 20 ;  
A. Kyatkin, *Phys. Lett.* **B361** (1995) 105;
- [9] R.Jackiw and S.Templeton, *Phys. Rev.* **D23** (1981) 2291; S.Deser, R.Jackiw and S.Templeton, *Phys. Rev. Lett.* **48** (1982) 975;*Ann. Phys. (NY)* **140** (1982) 372;
- [10] K.V.Zhukovsky and P.A.Eminov, *Izv. Vissht. Uch. Zav.* **5** (1995) 61-65;  
K.V.Zhukovskii and P.A. Eminov, *Phys. Lett.* **B359** (1995) 155;  
K.V.Zhukovsky and P.A.Eminov, *Yad. Fiz.* **59** (1996), 1265-1269;  
A.V.Borisov and K.V. Zhukovsky, *Vestn. Mosk. Univ., Fizika, Astron.*, **1** (1997) 74- 75
- [11] V.Ch.Zhukovsky, N.A.Peskov and A.Yu.Afinogenov, to appear in *Yad.Fiz.* **N5** (1998)
- [12] V. Zeitlin, e-Print Archive: hep-th/9612225, e-Print Archive: hep-th/9701100,
- [13] A.N.Sissakian, O.Yu.Shevchenko and S.B.Solganik, e-Print Archive: hep-th/9608159, e-Print Archive: hep-th/9612140
- [14] V.Ch.Zhukovskii and Sh.A.Mamedov, *Izv.Vissht.Uch.Zav., Fizika* **N10** (1990) 23
- [15] V.Ch.Zhukovskii and I.V.Mamsurov, to appear in *Yad.Fiz.* **N12** (1997)
- [16] S.M.Carroll, G.B.Field and R.Jackiw, *Phys. Rev.* **D 41** (1990) 1231;

- [17] D.Ebert and V.Ch.Zhukovsky, *Some Radiative and Topological Effects in Gauge Models with Vacuum Condensate*, in *Proceedings of the 31st International Symposium Ahrenshoop on the Theory of Elementary Particles*, Buckow, 2-6 September, 1997
- [18] V. Yu. Zeitlin, *Yad.Fiz.* **49** (1989) 742
- [19] A.N.Redlich, *Phys. Rev. Lett.* **52** (1983) 18;  
*Phys. Rev. D* **29** (1984) 2366
- [20] B. Nodland and J. P. Ralston, *Phys. Rev. Lett.*, **78**, (1997) 3047;  
*Phys. Rev. Lett.*, **79**, (1997) 1958.
- [21] S.M.Carroll and G.B.Field, *Phys. Rev. Lett.* **79** (1997) 2394; e-Print Archive: astro-ph/9704263;
- [22] J. P. Ralston and B. Nodland, e-Print Archive: astro-ph/9708114